



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore -\frac{4\theta^2}{\pi^2}\left(1+\frac{1}{3^2}+\frac{1}{5^2}+\dots\right)-\frac{4^2\theta^4}{2}\left(1+\frac{1}{3^4}+\frac{1}{5^4}+\dots\right)-\dots$$

$$-\frac{4^8\theta^{16}}{8\pi^{16}}\left(1+\frac{1}{3^{16}}+\frac{1}{5^{16}}+\frac{1}{7^{16}}+\dots\right)=-A-\frac{1}{2}A^2-\dots-\frac{1}{8}A^8-\dots$$

Equating like coefficients of θ^{16} we get

$$\frac{8192}{\pi^{16}}\left(1+\frac{1}{3^{16}}+\frac{1}{5^{16}}+\frac{1}{7^{16}}+\dots\right)=\frac{929569}{6435 \times 1260^2}.$$

$$\therefore \frac{6435}{2} \cdot \frac{161280^2}{929569} \left(1+\frac{1}{3^{16}}+\frac{1}{5^{16}}+\frac{1}{7^{16}}+\dots\right)=\pi^{16}.$$

Also solved by F. Anderegg, and G. W. Greenwood.

249. Proposed by J. J. KEYES, Fogg High School, Nashville, Tenn.

Solve $x+y+z=5$, $x^2+y^2=z^2$, $x^3+y^3+z^3=8$.

Solution by M. R. BECK, Cleveland, Ohio.

$x+y+z=5$, or $x+y=5-z$(1), $x^2+y^2=z^2$(2),

$x^3+y^3+z^3=8$, or $(x+y)(x^2-xy+y^2)=8-z^3$(3).

From (1) and (2), we have $xy=\frac{25-10z}{2}$(4).

Substituting (1), (2), and (4) in (3), and solving, $z=\frac{47}{5}$. Substituting $z=\frac{47}{5}$ in (1) and (2) and solving, $x=\frac{39}{5} \mp \frac{7}{5}\sqrt{-34}$, and $y=\frac{39}{5} \pm \frac{7}{5}\sqrt{-34}$.

Also solved by Henry Heaton, A. H. Holmes, L. E. Newcomb, J. Scheffer, and G. B. M. Zerr.

AVERAGE AND PROBABILITY.

172. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

What is the average length of all straight lines that can be drawn within a given triangle?

II. Solution by HENRY HEATON, Atlantic, Iowa.

Let ABC be the triangle. Let x =length of one of the straight lines, and θ the angle it would make with the side AB if produced to meet it, θ being taken $>$ than the angle A and $<$ than $\pi-B$. At the distance $x\sin\theta$ from AB draw ED parallel to AB cutting AC in E and BC in D . Then the number of lines of length x making the angle θ with AB is equal to the number of points in the triangle DEC whose area is $\frac{\Delta (b\sin A - x\sin\theta)^2}{b^2\sin^2 A}$. Similarly, the number of lines of length x making the angle θ with AC is equal to the number of points in the triangle whose area is $\frac{\Delta (c\sin A - x\sin\theta)^2}{c^2\sin^2 A}$, and the number making same angle with the side BC is equal to the number of points in the triangle whose area is